

Quality

Entire males – challenges and opportunities

Production and detection of intact boars and alternative processing routes for boar tainted products

By Ronald E. Klont, Esra Kurt, Lourens Heres and Bert Urlings

References

1. BABOL, J., SQUIRES, E.J. and K. LUNDSTRÖM (1996): Investigation of factors responsible for the development of boar taint. *Food Research International* 28, 573-581.
2. BANON, S., E. COSTA, M.D. GIL and M.-D. GARRIDO (2003): A comparative study of boar Saint in cooked and dry-cured meta. *Meat Sci.* 63, 381–88.
3. BONNEAU, M., KEMPSTER, A. J., CLAUS, R., CLAUDI-MAGNUSSEN, C., DIESTRE, A., TORNBORG, E., WALSTRA, P., CHEVILLON, P., WEILER, U. and G.L. COOK (2000): An international study on the importance of androstenone and skatole for boar taint: I. Presentation of the programme and measurement of boar taint compounds with different analytical procedures. *Meat Sci.* 54, 251–259.
4. BONNEAU, M., P. WALSTRA, C. CLAUDI-MAGNUSSEN, A.J. KEMPSTER, E. TORNBORG, K. FISCHER, A. DIESTRE, F. SIRET, P. CHEVILLON, R. CLAUS, G. DIJKSTERHUIS, P. PUNTER, K. R. MATTHEWS, H. AGERHEM, M. P. BÉAGUE, M. A. OLIVER, M. GISPERT, U. WEILER, G. VON SETH, H. LEASK, M. FONT I FURNOLS, D. B. HOMER and G. L. COOK (2000): An international study on the importance of androstenone and skatole for boar taint: IV. Simulation studies on consumer dissatisfaction with entire male pork and the effect of sorting carcasses on the slaughter line, main conclusions and recommendations. *Meat Sci.* 54, 285–295.
5. CLAUS, R., A. FISCHER and B. VOGELBACHER (1985): Konzentrationen des Ebergeruchsteroids im Schlachtkörper des Ebers un daraus hergesteller Fleischzeugnisse. *Fleischwirtsch.* 65 (3), 375–377.
6. CLAUS, R., WEILER, U., and A. HERZOG (1994): Physiological aspects of androstenone and skatole formation in the boar: a review with experimental data. *Meat sci.* 38, 289–305.
7. DIJKSTERHUIS, G.B., B. ENGEL, P. WALSTRA, M. FONT I FURNOLS, H. AGERHEM, K. FISCHER, M.A. OLIVER, C. CLAUDI-MAGNUSSEN, F. SIRET, M.P. BEAGUE, D.B. HOMER and M. BONNEAU (2000): An international study on the importance of androstenonen and skatole for boar taint: II. Sensory evaluation by trained panels in seven European countries. *Meat Sci.* 54, 261–269.
8. DUNSHEA, F.R., C. COLANTONI, K. HOWARD, I. MCCAULEY, P. JACKSON, K.A. LONG, S. LOPATICKI, E.A. NUGENT, J.A. SIMONS, J. WALKER and D.P. HENNESEY (2001): Vaccination of boars with GnRH vaccine (Improvac) and eliminates boar taint and increases growth performance. *Journal of Animal Sci.* 79, 2524–2535.
9. FONT I FURNOLS M., J. GONZÁLEZ, M GISPERT, M.A. OLIVER, M. HORTÓS, J. PÉREZ, P. SUÁREZ, L. and GUERRERO (2009): Sensory characterization of meat from pigs vaccinated against gonadotropin releasing factor compared to meat from surgically castrated, entire male and female pigs. *Meat Sci.* 83, 438–442.

10. GERRITZEN M.A., M. KLUIVERS-POODT, H. G. M. REIMERT, V. HINDLE and E. LAMBOOIJ (2008): Castration of piglets under CO₂-gas anaesthesia *Animal*, 2:11, pp 1666–1673.
11. GERRITZEN M.A., M. KLUIVERS-POODT, H. G. M. REIMERT, V. HINDLE and E. LAMBOOIJ (2009): Response to the Letter to the Editor on the surgical castration of piglets. *Animal*, 3:11, pp 1476–1477.
12. HAUGEN, J-E. (2003): Rapid detection of boar taint with chemical sensor technology. The challenge of gas-sensor based methods. European Association of Animal Production Working Group on Production and Utilisation of Meat from Entire Males, 13-14 November, Dublin, Ireland.
13. HUBER-EICHER, B. and P. SPRING (2008): Attitudes of Swiss consumers towards meat from entire or immunocastrated boars: A representative survey. *Research in Veterinary Sci.* 85, 626–627.
14. JOHNSON, L.A., RATH, D. VASQUES, J.M., MAXWELL and J.R. DOBRINSKY (2005): Preselection of sex of offspring in swine for production: current status of the process and its application. *Theriogenology*, 63, 615–624.
15. KLUIVERS-POODT, M. H. HOPSTER and H.A.M SPOOLDER (2007): Castration under anaesthesia and/or analgesia in commercial pig production. Report 85, Animal Science Group Lelystad, Wageningen-UR, The Netherlands.
16. LEE, G.J. A.L. ARCHIBALD, A.S. LAW, S. LLOYD, J. WOOD and C.S. HALEY (2005): Detection of quantitative trait loci for androstenone, skatole and boar taint between large white and Meishan pigs. *Animal Genetics* 36, 14–22.
17. LUNDE K., B. EGELANDSDAL, J. CHOINSKI, M. MIELNIK, A. FLATTEN and E. KUBBEROD (2008): Marinating as a technology to shift sensory thresholds in ready-to-eat entire male pork meat. *Meat Sci.* 80, 1264–1272.
18. MALMFORS, B. and I. HANSON (1974): Incidence of boar taint in Swedish Landrace and Yorkshire boars. *Livestock Production Sci.*, 1, 411–420.
19. MALMFORS, B. and K. LUNDSTRÖM (1983): Consumer reactions to boar taint, a review. *Livestock production Sci.* 10, 187–196
20. MATTHEWS, K. R., HOMER, D. B., PUNTER, P., BEAGUE, M.P., GISPERT, M., KEMPSTER, A.J., AGERHEM, H., CLAUDI-MAGNUSSEN, C., FISCHER, K., SIRET, F., LEASK, H., FONT I FURNOLS, M. and M. BONNEAU (2000): An international study on the importance of androstenone and skatole for boar taint: III. Consumer survey in seven European countries. *Meat Sci.* 54, 271–283.
21. MCCAULEY, I., D.P. HENNESEY, V. BOGHOSSION, L. SALI, L. SALVATORE, J. REYNOLDS and R. MAWSON (1997): Effects of cooking and processing pork on the perception of boar taint. In Bonneau, M, K. Lundstrom, & B. Malmfors (editors), *Boar taint in entire male pigs*. E.A.A.P Publication no. 92, Wageningen Pers.
22. MOE, M., T. MEUWISSEN, S. LIEN, C. BENDIXEN, X. WANG, L.N. CONLEY, I. BERGET, H. TAJET and E. GRINDFLEK (2007): Gene expression profiles in testis of pigs with extreme high and low levels of androstenone. *BMC Genomics* 8, doi:10.1186/1471-2164-8-405.
23. MOE, M., L. S. LIEN, T. AASMUNDSTAD, T. H.E. MEUWISSEN, M.H.S. HANSEN, C. BENDIXEN and E. GRINDFLEK (2009) Association between SNPs within candidate genes and compounds related to boar taint and reproduction *BMC Genetics* 2009, 10:32.
24. MORTENSEN, A.B. and S.E. SORENSEN (1984): Relationship between boar taint and skatole determined with a new analysis method. 30th European meeting of Meat Research Workers, Bristol, United Kingdom.

25. MOTTRAM, D.S., J.D. WOOD and R.L.S. PATTERSON (1982): Comparison of bulls and castrates for bacon production. 3. Composition and eating quality of bacon. *Animal Production* 35, 75-80.
26. MÜHLBAUER, I. W. OTTEN, W. LÜPPING, A. PALZER, S. ZÖLS, S. ELICKER, M. RITZMANN and K. HERINTITZL (2009): Investigation about CO₂ gas anaesthesia as an alternative option for piglet castration without anaesthesia. *Der Praktische Tierarzt* 90 (5) 460-464.
27. PAULY P., J.V. SPRING, O. DOHERTY, S. AMPUERO KRAGTEN and G. BEE (2008): Performances, meat quality and boar taint of castrates and entire male pigs fed a standard and a raw potato starch-enriched diet. *Animal* 2:11, 1707–1715.
28. QUINTANILLA, R., O. DEMEURE, J. P. BIDANEL, D. MILAN, N. IANNUCCELLI, Y. AMIGUES, J. GRUAND, C. RENARD, C. CHEVALET and M. BONNEAU (2003): Detection of quantitative trait loci for fat androstenone levels in pigs. *J. Anim. Sci.* 81, 385–394.
29. RATH, D. (2002): Low dose insemination in the sow. A review. *Reproduction in Domestic Animals*, 37, 201-205.
30. STOLZENBACH, S., G. LINDAHL, K. LUNDSTRÖM, G. CHEN and D.V. BYRNE (2009): Perceptual masking of boar taint in Swedish fermented sausages. *Meat Sci.* 81, 580–588.
31. SQUIRES, E.J. (2006): Possibilities for selection against boar taint. *Acta Veterinaria Scandinavica*, 48 (Suppl 1), S8.
32. TAJET, H., O. ANDRESEN and T.H.E. MEUWISSEN (2006): Estimation of genetic parameters of boar taint; skatole and androstenone and their correlations with sexual maturation. *Acta Veterinaria Scandinavica*, 48 (Suppl 1), S9.
33. TER BEEK, V. (2007) Sex sorting technology could save time and money. *Pig Progress*, 23 (10), 22–23.
34. VARONA, L., O. VIDAL, R. QUINTANILLA, M. GIL, A. SÁNCHEZ, J. M. FOLCH, M. HORTOS, M. A.RIUS, M. AMILLS and J. L. NOGUERA (2005): Bayesian analysis of quantitative trait loci for boar taint in a Landrace outbred population. *Journal of Animal Science* 83, 301–307.
35. WALSTRA, P. (1974): Fattening of young boars: quantification of negative and positive aspects. *Livestock Production Science*, 1, 187–196.
36. WALSTRA, P., C. CLAUDI-MAGNUSSEN, P. CHEVILLON, G. VON-SETH, A. DIESTRE, K.R. MATTHEWS, D.B. HOMER, and M. BONNEAU (1999): An international study on the importance of androstenone and skatole for boar taint: levels of androstenone and skatole by country and season. *Livestock Production Science*, 62, 15-28.
37. WEILER, U. M. FONT I FURNOLS, K. FISCHER, H. KEMMER, M.A. OLIVER, M. GISPERTT, A. DOBROWOLSKI and R. CLAUS (2000): Influence of differences in sensitivity of Spanish and German consumers to perceive androstenone on the acceptance of boar meat differing in skatole and androstenone concentrations. *Meat Sci.* 54, 297-304.
38. WILLEKE, H. (1993): Possibilities of breeding for low 5 α -androstenone content in pigs. *Pig News and Information*, 14, 31-33.
39. WILLIAMS, L. D., PEARSON, A. M. and N. B. WEBB (1963): Incidence of sex odor in boars, sows, barrows, and gilts. *Journal of Animal Sci.* 22, 116–168.
40. WRIGHT A.J., M. WHITING and A. TAYLOR (2009): Letter to the Editor on the surgical castration of piglets. *Animal*. 3 (11), 1474–1475.
41. XUE, J. L., DIAL, G.D., HOLTON, E.E., VICKERS, Z., SQUIRES, E.J., LOU, Y., GOTBOUT, D. and N. MOREL (1996): Breed differences in boar taint: relationship between tissue levels of boar taint compounds and sensory analysis. *Journal of Animal Sci.* 74, 2170–2177.

Authors' address

Dr. Ronald E. Klont, Dr. Esra Kurt, Dr. Lourens Heres and Prof. Dr. Bert Urlings, Vion Food Group, Boxtel, 5280 AA Netherlands, ronald.klont@vionfood.com